

A correspondence between H_3^+ WZW and Liouville theories on discs

Kazuo Hosomichi^a

^a*Service de Physique Théorique, CEA Saclay, F-91191 Gif-sur-Yvette Cedex, France*

We discuss how the disc correlators of H_3^+ WZW model are determined in terms of those of Liouville theory. (A presentation based on a collaboration with S. Ribault [1].)

Recently an interesting relation has been discovered between conformal blocks of affine $SL(2)$ and Virasoro algebras. By using it, the correlation functions of H_3^+ WZW model on sphere have been related to those in Liouville theory [2].

In this note we give a summary of the work [1] which generalized this correspondence to disc (or the upper half plane) and lead to an exact determination of disc correlators in H_3^+ .

We begin by reviewing the correspondence for sphere correlators. The space H_3^+ is a three dimensional hyperboloid in flat $\mathbb{R}^{3,1}$,

$$x_0^2 - x_1^2 - x_2^2 - x_3^2 = k\alpha' > 0, \quad x_0 > 0.$$

The space is coordinatized by a 2×2 hermite matrix h with unit determinant and positive trace. The CFT of interest is defined by the standard WZW action for $h(z, \bar{z})$ at level k , and has affine $SL(2)$ symmetry [3]. We define the components $J^a(z)$ and $\bar{J}^a(\bar{z})$ ($a = \pm, 3$) of the currents $-k\partial h h^{-1}$, $kh^{-1}\bar{\partial}h$ so that they are hermite conjugate to each other, and consider correlators of “ μ -basis” primary fields $\Phi_j(\mu|z)$ which have

$$L_0 = \frac{-j(j+1)}{k-2}, \quad J_0^- = i\mu, \quad \bar{J}_0^- = i\bar{\mu}.$$

Their correlators satisfy the so-called KZ equations [4] that follow from the relation between the stress tensor and the currents.

We relate this model to Liouville theory with coupling $b = (k-2)^{-\frac{1}{2}}$ and central charge

$$c = 1 + 6Q^2, \quad Q \equiv b + 1/b.$$

The primary field $V_\alpha(z)$ with Liouville momentum α has conformal weight $\alpha(Q - \alpha)$. The operator with $\alpha = -\frac{1}{2b}$ corresponds to a degenerate

representation of Virasoro algebra, and correlators containing $V_{-1/2b}(z)$ are known to satisfy the so-called BPZ equation[5] which is a differential equation of second order in z .

An interesting relation [6] between KZ and BPZ equations leads to a correspondence between n -point blocks of affine $SL(2)$ algebra and $(2n-2)$ -point Virasoro conformal blocks. n of their insertions are related by $\alpha_a = b(j_a + 1) + \frac{1}{2b}$, and the additional $(n-2)$ insertions on Virasoro side are all with $\alpha = -\frac{1}{2b}$. This was used in [2] to relate the correlators of the two theories on sphere,

$$\left\langle \prod_{a=1}^n \Phi_{j_a}(\mu_a|z_a) \right\rangle = \delta^2(\sum_{a=1}^n \mu_a) \cdot |\Theta|^{\frac{1}{b^2}} |u|^2 \times \left\langle \prod_{a=1}^n V_{\alpha_a}(z_a) \prod_{a=1}^{n-2} V_{-\frac{1}{2b}}(y_a) \right\rangle. \quad (1)$$

Here u , y_a and Θ are (implicitly) given by

$$\sum_{a=1}^n \frac{\mu_a}{\zeta - z_a} = u \frac{\prod_{a=1}^{n-2} (\zeta - y_a)}{\prod_{a=1}^n (\zeta - z_a)},$$

$$\Theta = \prod_{a < \bar{a}} (z_a - z_{\bar{a}}) \prod_{a < \bar{a}} (y_a - y_{\bar{a}}) \prod_{a, \bar{a}} (z_a - y_{\bar{a}})^{-1}.$$

Once verified for $n \leq 3$, the correspondence (1) for larger n simply follows from the fact that the two sides behave in the same manner under factorizations $n \rightarrow n_1 + n_2$.

We wish to generalize this correspondence to correlators on discs. Though there are several types of D-branes in H_3^+ WZW model known (see e.g. [7]), here we focus on the one parameter family defined by a plane

$$x_3 = \sqrt{k\alpha'} \sinh r, \quad (2)$$

intersecting the hyperboloid. Our convention is such that they are all characterized by the boundary condition on currents, $J^3 = \bar{J}^3$, $J^\pm = \bar{J}^\pm$. The system therefore has an unbroken affine $SL(2, \mathbb{R})$ symmetry, and its zero mode part generates the isometry of the D-brane worldvolume H_2^+ . We denote by $\Psi_l(\nu|w)$ the boundary operator between D-branes (2) with eigenvalues

$$L_0 = -\frac{l(l+1)}{k-2}, \quad J_0^- = i\nu.$$

In Liouville theory, we consider boundary operators B_β with $L_0 = \beta(Q - \beta)$ and FZZT boundary conditions [8] labelled by s .

To formulate the correspondence between disc correlators, we first recall the relation between one-point functions found in [9],

$$\langle \Phi_j(\mu|z) \rangle_r \sim \langle V_\alpha(z) \rangle_{s=\frac{r}{2\pi b} - \frac{i}{2b} \text{sgn}(\text{Im}\mu)},$$

telling us how to relate the boundary conditions in two theories. Now the correspondence between general disc correlators should take the form

$$\begin{aligned} & \left\langle \prod_{a=1}^n \Phi_{j_a}(\mu_a|z_a) \prod_{a=1}^m \Psi_{l_a}(\nu_a|w_a) \right\rangle \\ &= \delta(\sum_a(\mu_a + \bar{\mu}_a) + \sum_a \nu_a) |u| \Theta^{\frac{1}{2b^2}} \cdot \\ & \quad \times \left\langle \prod_{a=1}^n V_{\alpha_a}(z_a) \prod_{a=1}^m B_{\beta_a}(w_a) \right. \\ & \quad \left. \prod_{a=1}^{n'} V_{-\frac{1}{2b}}(y_a) \prod_{a=1}^{m'} B_{-\frac{1}{2b}}(v_a) \right\rangle, \end{aligned} \quad (3)$$

where $2n' + m' = 2n + m - 2$, and (u, y_a, v_a) are determined via a real function of $\zeta \in \mathbb{R}$,

$$\begin{aligned} \varphi(\zeta) &\equiv \sum_{a=1}^n \left(\frac{\mu_a}{\zeta - z_a} + \frac{\bar{\mu}_a}{\zeta - \bar{z}_a} \right) + \sum_{a=1}^m \frac{\nu_a}{\zeta - w_a} \\ &= u \cdot \frac{\prod_{a=1}^{n'} |\zeta - y_a|^2 \prod_{a=1}^{m'} (\zeta - v_a)}{\prod_{a=1}^n |\zeta - z_a|^2 \prod_{a=1}^m (\zeta - w_a)}. \end{aligned}$$

Θ is a function of coordinate differences similar to the case of sphere. The most important part of the correspondence is how the boundary conditions in two theories are related. The boundary of the disc is cut at $\{w_a\}$ into segments labelled

by different r , and is cut further at $\{v_a\}$ in Liouville theory side. Then the rule is that *a boundary segment labelled by r in H_3^+ model maps to a Liouville FZZT brane with*

$$s = \frac{r}{2\pi b} - \frac{i}{4b} \text{sgn}\varphi(\zeta).$$

Obviously s is constant on each segment, as it should. Our rule is also consistent with the fact in Liouville theory that s has to jump by $\pm \frac{i}{2b}$ where $B_{-1/2b}$ is inserted [8].

It is easy to show that our formula (3) behaves consistently under different factorization limits, which actually serves as a strong consistency check of our formula. On the other hand, different factorization limits of a correlator of H_3^+ model are usually not smoothly connected because of transmutations of degenerate operators in Liouville theory along the way:

$$V_{-1/2b}(y_a) \longleftrightarrow B_{-1/2b}(v_a) B_{-1/2b}(v_{a'}).$$

The construction of disc higher point functions from basic ones is therefore not so straightforward in H_3^+ WZW model. Our work [1] has argued that the disc correlators must be continuous at transmutation points, and has shown that the formula (3) is very likely the only solution to the continuity as well as other consistency requirements.

The formula (3) has been used in [1] to compute some disc structure constants of H_3^+ WZW model which were not known before. We have got the two point function of boundary operators which is fully consistent with that of $N = 2$ Liouville theory [10], and the bulk boundary propagator with the correct semiclassical behavior at large k . These serve as another non-trivial check of our formula.

REFERENCES

1. K. Hosomichi and S. Ribault, hep-th/0610117.
2. S. Ribault et.al., JHEP **0506** (2005) 014.
3. K. Gawedzki, Nucl. Phys. B**328** (1989) 733.
4. V.G. Knizhnik et.al., Nucl. Phys. B**247** (1984) 83.
5. A.A. Belavin et.al., Nucl. Phys. B**241** (1984) 333.
6. A.V. Stoyanovsky, math-ph/0012013.
7. B. Ponsot et.al., JHEP **0202** (2002) 016.
8. V. Fateev et.al., hep-th/0001012.
9. S. Ribault, JHEP **0608** (2006) 015.
10. K. Hosomichi, hep-th/0408172.